

OPIC

OFFICE DE LA PROPRIÉTÉ  
INTELLECTUELLE DU CANADA

CIPO

CANADIAN INTELLECTUAL  
PROPERTY OFFICE

REC'D 26 MAY 1999

WIPO PCT

09/647965

*Bureau canadien  
des brevets**Canadian Patent  
Office* 5*Certification*

La présente atteste que les documents  
ci-joints, dont la liste figure ci-dessous,  
sont des copies authentiques des docu-  
ments déposés au Bureau des brevets.

*Certification*

This is to certify that the documents  
attached hereto and identified below are  
true copies of the documents on file in  
the Patent Office.

Specification and Drawings, as originally filed, with Application for Patent Serial No:  
2,234,588, on April 7, 1998, by THE SIR MORTIMER B. DAVIS - JEWISH  
GENERAL HOSPITAL, assignee of John Hiscott and Rongtuan Lin, for "Highly Active  
Forms of Interferon".

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)



Agent certificateur / Certifying Officer

May 19, 1999

Date

Industrie  
CanadaIndustry  
Canada

(CIPO 68)

Canada

VIRUS INDUCED PHOSPHORYLATION OF INTERFERON REGULATORY  
FACTOR-3

BACKGROUND OF THE INVENTION

Interferons (IFNs) are a large family of multifunctional secreted proteins involved in antiviral defence, cell growth regulation and immune activation (63). Virus infection induces the transcription and synthesis of multiple IFN genes (33,52,63); newly synthesized IFN interacts with neighbouring cells through cell surface receptors and the JAK-STAT signalling pathway, resulting in the induction of over 30 new cellular proteins that mediate the diverse functions of the IFNs (17,35,39,58). Among the many virus- and IFN-inducible proteins are the growing family of IRF transcription factors, the Interferon Regulatory Factors (IRFs).

IRF-1 and IRF-2 are the best characterized members of this family, originally identified by studies of the transcriptional regulation of the human IFN- $\beta$  gene (22,23,30,47). Their discovery preceded the recent expansion of this group of IFN-responsive proteins which now include seven other members: IRF-3, IRF-4/Pip/ICSAT, IRF-5, IRF-6, IRF-7, ISGF3 $\gamma$ /p48 and ICSBP (48). Structurally, the Myb oncoproteins share homology with the IRF family, although its relationship to the IFN system is unclear (62). Recent evidence also demonstrates the presence of virally encoded analogue of cellular IRFs - vIRF in the genome of human herpes virus 8 (HHV8) (55).

The presence of IRF-like binding sites in the promoter region of the IFNA and IFNB genes implicated the IRF factors as essential mediators of the induction of IFN genes. The original results of Harada et al. (30,32) indicated that IFN gene induction was activated by IRF-1, while the related IRF-2 factor suppressed IFN expression. However, the essential role of IRF-1 and IRF-2 in the regulation of IFNA and IFNB gene expression has become controversial with the observation that mice containing homozygous deletion of IRF-1 or IRF-2, or fibroblasts derived from these mice, induced IFNA and IFNB gene expression after virus infection to the same level as the wild-type mice or cells (44).

On the other hand, IRF-1 was shown to have an important role in the antiviral effects of IFNs (44,54). IRF-1 binds to the ISRE element present in many IFN-inducible gene promoters and activates expression of some of these genes (54). However, activation of ISG genes by IFNA and IFNB was shown to be mediated generally by the multiprotein ISGF3 complex (31,36,38). The binding of this complex to DNA is mediated by another member of the IRF family, ISGF3 $\gamma$ /p48, which in IFN-treated cells interacts with phosphorylated STAT1 and STAT2 transcription factors forming the heterotrimeric complex ISGF3 (8,39,62). The homozygous deletion of p48 in mice abolished the sensitivity of these mice to the antiviral effects of IFNs, and virus-infected macrophages from p48 $^{-/-}$  mice showed an impaired induction of IFNA and IFNB genes (31).

Several other members of the IRF family have been identified. The ICSBP gene is expressed exclusively in the cells of the immune system (18,64) and its expression can be enhanced by IFN $\gamma$ . ICSBP was shown to form a complex with IRF-1 and inhibit the transactivating activity of IRF-1 (9,59). The homozygous deletion of ICSBP in mice leads to defects in myeloid cell lineage development and chronic myelogenous leukemia (34). Another lymphoid specific Pip/LSIRF/IRF-4 was identified (19,43,66) that interacts with phosphorylated PU.1, a member of the Ets family of transcription factors (15). The Pip/PU.1 heterodimer can bind to the immunoglobulin light chain enhancer and function as a B cell specific transcriptional activator. Expression of Pip/LSIRF was induced by antigenic stimulation but not by IFN, and Pip/LSIRF/IRF-4 -/- mice failed to develop mature T and B cells (46). A novel member of the IRF family was recently identified by its ability to bind to an ISRE-like element in the promoter region of the Qp gene of EBV (69).

Another unique member of the human IRF family, IRF-3 was characterized recently (2). The IRF-3 gene encodes a 55-kDa protein which is expressed constitutively in all tissues. Recombinant IRF-3 binds to the ISRE element of the IFN-induced gene ISG-15 and stimulates this promoter in transient expression assays. In contrast to IRF-1, which contains both DNA binding and transactivating domains, IRF-3 - like IRF-2 and ICSBP - does not contain a well-defined transactivation domain. Viral induction or IFN-treatment does not stimulate the expression of the IRF-3 gene. In

previous studies, applicant showed that IRF-3 binds to the IE and PRDIII regions of the IFNA and IFNB promoters respectively, but has different effects on their transcriptional activity (56). While the induction of the

IFNA4 promoter activated by IRF-1 or virus infection was inhibited in the presence of IRF-3, the fusion protein containing the IRF-3 DNA binding domain and the RelA(p65) transactivation domain effectively activated both IFNA and IFNB promoters. In contrast, co-expression of IRF-3 and RelA plasmids transactivated the IFNB gene promoter, but not the promoter of the IFNA4 gene (56).

The applicant has examined the virus-induced, post-translational modulation of IRF-3 protein following Sendai virus infection. Applicant's studies demonstrate that phosphorylation represents an important post-translational modification of IRF-3 leading to cytoplasmic to nuclear translocation of phosphorylated IRF-3, stimulation of DNA binding and transcriptional activity, association of IRF-3 with the transcriptional co-activator CBP/p300, and ultimately proteasome mediated degradation.

#### SUMMARY OF THE INVENTION

The interferon regulatory factors (IRF) consist of a growing family of related transcription proteins first identified as regulators of the IFN $\alpha/\beta$  gene promoters, as well as the ISRE of some IFN-stimulated genes. IRF-3 was originally identified as a member of the IRF family based on homology with other IRF family members and on binding to the ISRE of the ISG15 promoter. IRF-3 is expressed

~~constitutively in a variety of tissues and the relative~~  
levels of IRF-3 mRNA do not change in virus-infected or  
IFN-treated cells.

Applicant has demonstrated that, following Sendai  
virus infection, IRF-3 is post-translationally modified by  
protein phosphorylation at multiple serine and threonine  
residues, located in the carboxy-terminus of IRF-3. A  
combination of IRF-3 deletion and point mutations localized  
the inducible phosphorylation sites to the region  
10 -ISNSHPLSLTSDQ- between amino acids 395 and 407; point  
mutation of Ser-396 and Ser-398 residues eliminated  
virus-induced phosphorylation of IRF-3 protein, although  
residues Ser-402, Thr-404 and Ser-405 were also targets.  
Phosphorylation results in the cytoplasmic to nuclear  
translocation of IRF-3, DNA binding and increased  
transcriptional activation. Substitution of the Ser/Thr  
sites with the phosphomimetic Asp generated a constitutively  
active form of IRF-3 that functioned as a very strong  
activator of promoters containing PRDI/PRDIII or ISRE  
20 regulatory elements. Use of phosphomimetic Glu for this  
purpose is also possible. Phosphorylation also appears to  
represent a signal for virus mediated degradation, since the  
virus induced turnover of IRF-3 was prevented by mutation of  
the IRF-3 Ser/Thr cluster or by proteasome inhibitors.

Interestingly, virus infection resulted in the  
association of IRF-3 with the CBP coactivator, as detected by  
co-immunoprecipitation with anti-CBP antibody, an interaction  
mediated by the C-terminal domains of both proteins.

Mutation of the residues Ser-396 and Ser-398 in IRF-3 abrogated its binding to CBP. These results are discussed in terms of a model in which virus-inducible C-terminal phosphorylation of IRF-3 alters protein conformation to permit nuclear translocation, association with transcriptional partners and primary activation of IFN- and IFN-responsive genes.

#### DETAILED DESCRIPTION OF THE INVENTION

Several novel aspects of post-translational regulation of the IRF-3 transcription factor are highlighted in the present study. 1) Sendai virus dependent phosphorylation of IRF-3 was detected, occurring in a cluster of Ser and Thr sites in the carboxyl-terminal end of the protein. The residues implicated in this regulatory phosphorylation event are Ser-396/Ser-398/Ser-402/Thr-404/Ser-405, particularly the Ser-396/Ser-398 amino acids. 2) Phosphorylation of the IRF-3 in the Ser-Thr cluster resulted in the cytoplasmic to nuclear translocation of IRF-3; nuclear translocation was blocked by mutation of the phosphorylated amino acids. 3) Sendai virus infection induced the DNA binding and transactivation potential of IRF-3. Furthermore, IRF-3 containing the phosphomimetic Asp at the sites of C-terminal phosphorylation was an exceptionally strong transactivator of PRDI/PRDIII and ISRE containing promoters. 4) Phosphorylation was also required for the association of IRF-3 with the CBP co-activator protein. 5) Sendai virus infection resulted in IRF-3 degradation; again, phosphorylation was required as a

signal for inducer-mediated degradation since mutation of Ser/Thr cluster also blocked virus induced degradation.

Cytoplasmic to nuclear translocation of IRF-3 as a consequence of virus infection was inhibited by mutation of the Ser/Thr cluster, indicating an important regulatory role for C-terminal phosphorylation in the activation of IRF-3. Also strikingly, the conversion of the phosphorylation sites to the phosphomimetic Asp altered the subcellular localization of IRF-3 in uninfected cells. A proportion of IRF-3(5D) was localized to the nucleus of uninfected cells, suggesting that some IRF-3 may shuttle to and from the nucleus constitutively; this observation is consistent with the identification of a nuclear export signal in IRF-3. Mutation of L144A/L145A in the NES element produced the most impressive alterations in subcellular localization. In uninfected cells, IRF-3 was partitioned in both the nucleus and cytoplasm; virus infection changed the nuclear pattern of staining from extra-nucleolar homogeneous staining as observed for wtIRF-3 to an intense nuclear speckling. At this stage, the nature of the subnuclear changes in IRF-3 localization are not explained, although it is possible that IRF-3(NES) translocates efficiently into the nucleus but becomes trapped in the nuclear pore complex during the export process.

One of the striking results of the mutagenesis of the C-terminal domain of IRF-3 was the generation of IRF-3(5D), an exceptionally strong activator of IFN- $\beta$  and ISG-15 gene expression. The phosphomimetic form of IRF-3



—alone—was—able—to stimulate IFN- $\beta$  expression as strongly as virus infection, a level of stimulation not previously observed in co-expression experiments (24,61). In previous experiments, applicant demonstrated that IRF-3 was able to bind the ISRE element of ISG-15, as well as the PRDIII/PRDI and IE regions of the IFNB and IFNA promoters, respectively (2,56). Virus induction results in the appearance of two new protein-DNA complexes; supershift experiments confirmed that both complexes contain IRF-3; it is not clear at this stage whether the upper complex also contains other proteins such as in the VIC (10,29) and DRAF (16) complexes or whether the lower complex represents a breakdown product of IRF-3. Strikingly, the same complexes appeared following co-transfection of IRF-3(5D) expression plasmid in the absence of virus induction, indicating that IRF-3(5D) represented a constitutive DNA binding form of IRF-3. Thus, in uninfected cells, IRF-3(5D) localized in part to the nucleus (Fig. 5), interacted with DNA constitutively and was a strong activator of gene expression (Fig. 6).

The recent crystal structure of the related IRF-1 protein bound to PRDI provides evidence for a novel helix-turn-helix motif that latches onto a GAAA core sequence via three of the five conserved tryptophan amino acids of the DNA binding domain (20). By analogy with IRF-3, two GAAANN sequences present in PRDIII of IFN- $\beta$  and another GAAANN element present in PRDI may serve as DNA contacts for multiple IRF-3(5D) proteins with strong activating potential. Similarly, the ISRE element of the ISG-15 promoter also

~~contains several GAAANN anchors for potential IRF binding.~~

Given the range of promoters that possess this hexameric sequence (48), it will be of interest to determine the capacity of IRF-3(5D) to stimulate expression of different cytokine and chemokine genes.

IRF-3 joins a growing list of cellular and viral proteins that functionally interact with CBP/p300 proteins, highly homologous proteins originally identified through their interactions with adenovirus E1A and CREB proteins (1,13). As a critical determinant of its global transcriptional coactivator activity, CBP/p300 possesses histone acetyltransferase activity (5,50). Acetylation of histones is involved in the destabilization and remodelling of nucleosomes, a crucial step in permitting the accessibility of transcriptional factors to DNA templates. Several studies have now demonstrated that CBP/p300 participates in the transcriptional process by providing a scaffold for different classes of transcriptional regulators on specific chromatin domains (12,50). A growing body of biochemical and genetic evidence also implicates CBP/p300 as a negative regulator of cell growth, based on its interactions with adenovirus E1a, SV40 large T antigen and the tumour suppressor p53, among others. With regard to p53-CBP/p300 complex formation, functional interaction between these two important growth regulatory proteins accounts for several of the known activities of p53 (3,28,40); interestingly, CBP/p300 was shown recently to acetylate p53 and stimulate its transactivation potential

(27).

It will be of interest to determine whether IRF-3 is similarly modified by CBP association. The functional consequences of IRF-3 interaction with CBP/p300 remain to be elucidated, although recent studies demonstrated that CBP/p300 also functionally interacts with STAT 1 (68) and STAT 2 (7) and may contribute to IFN $\alpha$  and IFN $\gamma$  nuclear signalling. Recently published studies have demonstrated that synergistic activation of the IFN $\beta$  promoter requires recruitment of CBP/p300 to the enhanceosome, via a new activating surface assembled from the activation domains of all the transcription factors in the enhanceosome (37,45). Alterations in any of the activation domains decreased both CBP recruitment and transcriptional synergy. By analogy, recruitment of CBP/p300 to DNA bound IRF-3 is likely required for maximal transcriptional activation. Association requires the interaction of the C-terminal domain of IRF-3 and the C-terminal interaction domain of CBP, a region previously shown to associate with the p53 tumour suppressor, whereas STAT1 and STAT2 associate with different regions of CBP (7,68).

Virus induced phosphorylation of IRF-3 also represents a signal for proteasome mediated degradation of IRF-3, since mutation of the Ser-396/Ser-398 or the use of proteasome inhibitors prevented the post infection degradation of IRF-3. Virus induced degradation of IRF-3 is reminiscent of the virus-induced turnover of another member of the IRF family - IRF-2. In response to dsRNA or viral

induction, the 50 kD IRF-2 protein is proteolytically processed into a smaller, 24-27 kDa protein (51) comprising the 160 aa DBD of IRF-2, termed TH3 (14) or In4 (65).

Although TH3 has been shown to bind DNA and repress transcription more efficiently than the full length IRF-2 protein (42), its physiological role is not clear. Since the induction kinetics of TH3 are slower than that of IFN- $\beta$  in response to dsRNA or viral infection (14), it has been suggested that the IRF-2 cleavage product may be a post-induction repressor of IFN- $\beta$  gene expression (65).

Virus induced phosphorylation of IRF-3 at the C-terminal Ser/Thr residues and its subsequent degradation by a proteasome dependent pathway are also similar to the well studied phosphorylation and degradation of I $\kappa$ B $\alpha$  which leads to activation of NF- $\kappa$ B binding activity (reviewed in 4,6). In unstimulated cells, NF- $\kappa$ B heterodimers are retained in the cytoplasm by inhibitory I $\kappa$ B proteins. Upon stimulation by many activating agents, including cytokines, viruses and dsRNA, I $\kappa$ B $\alpha$  is rapidly phosphorylated and degraded, resulting in the release and nuclear translocation of NF- $\kappa$ B. The amino-terminus of I $\kappa$ B $\alpha$  represents a signal response domain for activation of NF- $\kappa$ B and substitution of alanine for either Ser-32 or Ser-36 completely abolished the signal-induced phosphorylation and degradation of I $\kappa$ B $\alpha$ , and blocked the activation of NF- $\kappa$ B. These mutations also blocked *in vitro* ubiquitination of the I $\kappa$ B $\alpha$  protein. The amino-terminus of I $\kappa$ B $\alpha$  is necessary for signal-induced phosphorylation and ubiquitination, but for degradation to

~~occur, there is an absolute requirement for the C-terminal~~  
PEST domain (reviewed in 4,6).

Similarities and differences exist between the  
observed degradation of IRF-3 and the mechanism of I $\kappa$ B $\alpha$   
degradation. The C-terminal phosphorylation of IRF-3 as a  
consequence of virus infection is required for its subsequent  
degradation based on the deletion and point mutation analysis  
of the region -ISNSHPLSLTSDQ- between amino acids 395 and  
407. Minimally, phosphorylation of Ser-396 and Ser-398 are  
10 required for subsequent degradation, although Ser-402,  
Ser-404 and Ser-405 may represent secondary phosphorylation  
sites. Likewise, in the case of I $\kappa$ B $\alpha$ , phosphorylation and  
Ser-32 and Ser-36 are required for inducer mediated  
degradation. Furthermore, the protease inhibitor calpain  
inhibitor I and the more specific proteasome inhibitor MG132  
block IRF-3 turnover.

A major difference in the mechanisms of I $\kappa$ B $\alpha$  and  
IRF-3 turnover lies in the nature of the inducing stimuli.  
Multiple inducers - cytokines such as TNF and IL-1, viruses,  
20 LPS, oxidative stress, etc (6) - all lead to the induction of  
I $\kappa$ B $\alpha$  phosphorylation and degradation whereas IRF-3  
phosphorylation appears to be induced only by virus infection  
and dsRNA addition; other inducers have not resulted in IRF-3  
turnover.

A significant temporal difference also exists  
between I $\kappa$ B $\alpha$  phosphorylation/turnover and IRF-3  
phosphorylation/degradation. Many activators of NF- $\kappa$ B  
stimulate I $\kappa$ B $\alpha$  phosphorylation within minutes and TNF induced

degradation occurs within the first 15-30 min after treatment. In the case of IRF-3, phosphorylation is not detected until 6-8 hours after infection and continues in a heterogenous manner over the next 10-12 hours. Previous experiments have, however, demonstrated that Sendai virus-induced turnover of I $\kappa$ B $\alpha$  also occurs slowly over several hours (24).

Based on the data presented herein and by analogy with the properties of other IRF family members (48), applicant proposes the following model to explain several observations. IRF-3 exists in a latent state in the cytoplasm of uninfected cells; the C-terminus may physically interact with the DNA binding domain in such a way as to obscure both the DBD and the IAD regions of the protein; the presence of an autoinhibitory domain within the C-terminal 20aa (407-427) would explain the activating effect of this deletion, as seen previously with IRF-4 (11,19). Virus induced phosphorylation at the Ser/Thr at 396-405aa cluster leads to a conformational change in IRF-3, exposing both the DBD and IAD and relieving C-terminal autoinhibition. Translocation to the nucleus, occurring via an unidentified nuclear localization sequence or in conjunction with a transcriptional partner associating through the IAD region, leads to DNA binding at ISRE- and PRDI/PRDIII-containing promoters. Phosphorylation is also necessary for IRF-3 association with the chromatin remodelling activity of CBP/p300. The presence of a NES element ultimately shuttles IRF-3 from the nucleus and terminates the initial activation

of IFN responsive promoters. The phosphorylated form of IRF-3 exported from the nucleus may now be susceptible to proteasome mediated degradation. This scenario shares several features with the protein synthesis independent activation of NF- $\kappa$ B, and further suggests that IRF-3 may represent a component of virus- or dsRNA-inducible complexes such as DRAF (16) or VIC (10,29) that could play a primary role in the induction of IFN- or IFN responsive genes.

#### DESCRIPTION OF THE FIGURES

10                   Figure 1. Sendai virus infection induces IRF-3 degradation.

IRF-3 expression plasmid CMVBL-IRF3 (lanes 1 and 2) or CMVBL vector alone (lanes 3 and 4), both at 5  $\mu$ g were transiently transfected into 293 cells by the calcium phosphate method. At 24h post transfection, cells were infected with Sendai virus for 16h (lanes 2 and 4) or left uninfected (lanes 1 and 3). Whole cell extracts (20  $\mu$ g) were prepared and analyzed by immunoblotting with anti-IRF-3 antibody.

20                   Figure 2. Sendai virus induced phosphorylation and degradation of IRF-3 protein.

A) rtTA-IRF-3 cells, selected as described in the Example, were induced to express IRF-3 by doxycycline treatment for 24h. At 24h after Dox addition, cells were infected with Sendai virus for 4, 8, 12, 16, 20, or 24h (lanes 2-7) or were left uninfected (lane 1). IRF-3 protein was detected in whole cell extracts (10  $\mu$ g) by immunoblot. Two forms of IRF-3 were detected, designated as form I and form II.

B) At 24h post Dox induction, rtTA-IRF-3 cells were infected

with Sendai virus for 16 hours (lanes 4-8) or were left uninfected (lanes 1-3). Whole cell extracts from untreated cells (20  $\mu$ g) or Sendai virus infected cells (60  $\mu$ g) were incubated with 0.3 units of potato acidic phosphatase (PPA, lanes 2, 3, 7 and 8) or 5 units of calf intestinal alkaline phosphatase (CIP, lanes 4 and 5) in the absence (lanes 1, 2, 4, 6 and 7) or presence of phosphatase inhibitors (lanes 3, 5 and 8). Phosphorylated IRF-3 protein appears as a distinct band in immunoblots, migrating more slowly than IRF-3 forms I and II.

Figure 3. Analysis of IRF-3 deletion mutants in Sendai virus induced phosphorylation.

(A) Schematic representation of four IRF-3 deletions. Thick solid lines and thin dashed lines indicate included and excluded sequences, respectively. The N-terminal IRF homology domain, the nuclear export signal (NES) and C-terminal IRF association domain are indicated.

(B) Expression plasmids (5  $\mu$ g each) encoding wild type and deletion mutants of IRF-3 (as indicated above the lanes) were transiently transfected into 293 cells; at 24h post transfection, cells were infected with Sendai virus for 16h (lanes 2, 4, 6, 8, and 10) or left uninfected (lanes 1, 3, 5, 7, and 9). Whole cell extracts (20  $\mu$ g) were prepared from infected and control cells and analyzed by immunoblotting for IRF-3 forms I and II and for the presence of phosphorylated IRF-3 (P-IRF-3) with anti-IRF-3 antibody.

Figure 4. Analysis of IRF-3 point mutations in Sendai virus induced phosphorylation.



~~(A) Schematic representation of IRF-3 point mutations.~~

Thick solid lines and thin dashed lines indicate included and excluded sequences, respectively. The N-terminal IRF homology domain, the Nes element and C-terminal IRF

association domain are indicated. Amino acids residues from 382 to 414 and from 141 to 147 are shown. The amino acids targeted for alanine or aspartic acid substitution are shown in large print. The point mutations are indicated below the sequence: (2A: S396A/S398A; 3A: S402A/T404A/S405A; 5A:

10 S396A/S398A/S402A/T404A/S405A); 5D  
S396D/S398D/S402D/T404D/S405D; J2A: S385A/S386A; NES:  
S145A/S146A).

(B) Expression plasmids (5  $\mu$ g each) encoding wild type and point mutants of IRF-3 (as indicated above the lanes) were transiently transfected into 293 cells; at 24h post transfection, cells were infected with Sendai virus for 16h (lanes 2, 4, 6, 8, 10, 12, 14, 16 and 18) or left uninfected (lanes 1, 3, 5, 7, 9, 11, 13, 15 and 17). Whole cell extracts (20  $\mu$ g) were prepared from infected and control  
20 cells and analyzed by immunoblotting for IRF-3 forms I and II and for the presence of phosphorylated IRF-3 (P-IRF-3) with anti-IRF-3 antibody.

Figure 5. Virus dependent cytoplasmic-nuclear translocation of IRF-3.

The subcellular localization of the GFP-IRF-3 (A and B), GFP-IRF-3(5A) (C and D), GFP-IRF-3(5D) (E and F) and GFP-IRF-3(NES) (G and H) was analyzed in uninfected (A, C, E, and G) and Sendai virus infected COS-7 cells at 16h after

infection. GFP fluorescence was analyzed in living cells with a Leica fluorescence microscope using 40x objective.

Figure 6. Transactivation of PRDI/PRDIII and ISRE containing promoters by IRF-3.

293 cells were transfected with IFN $\beta$ -CAT (A and B) or ISG15-CAT (C) reporter plasmids and the various expression plasmids as indicated below the bar graph. CAT activity was analyzed at 48h post-transfection with 100  $\mu$ g (IFN $\beta$ -CAT) or 10  $\mu$ g (ISG15-CAT) of total protein extract for 1-2h at 37°C.

Relative CAT activity was measured as fold activation (relative to the basal level of reporter gene in the presence of CMV-B1 vector alone after normalization with co-transfected  $\beta$ -Gal activity); the values represent the average of three experiments with variability shown in the error bar.

Figure 7. Stabilization of IRF-3 by proteasome inhibitors.

IRF-3  $\Delta$ N ( $\Delta$ 9-133) (B) or IRF-3  $\Delta$ N2A (C) expression plasmids were transiently transfected into 293 cells; at 24h post transfection, cells were infected with Sendai virus and treated for 12h with calpain inhibitor I (100  $\mu$ M, lanes 2 and 5) or MG132 proteasome inhibitor (40  $\mu$ M, lanes 3 and 6). Ethanol, the solvent for calpain inhibitor I and MG132, was added to the cells as control (lanes 1 and 4). Endogenous (A) and transfected (B and C) IRF-3 proteins were detected in whole cell extracts (20  $\mu$ g) by immunoblot.

Figure 8. IRF-3 interacts with CBP in virus infected cells.

(A) Schematic representation of CBP, illustrating the domains involved in interaction with host or viral proteins (modified from (28)) and the myc-tagged CBP proteins (CBP1, CBP2, CBP3) used for immunoprecipitation.

---

10 (B) 293 cells were transfected with wild type and deletion mutants of IRF-3 expression plasmid (5  $\mu$ g, as indicated above the lanes) or left untransfected (lanes 1 and 8). At 24h after transfection, cells were infected with Sendai virus for 16h (lanes 1, 3-8, and 10-13) or left uninfected (lanes 1 and 9). Whole cell extracts (300  $\mu$ g, except lane 1, which was 600  $\mu$ g) were immunoprecipitated with anti-CBP antibody A22 (lanes 1-6) or with preimmune serum (lane 7). The immunoprecipitated complexes (lanes 1-7) or 30  $\mu$ g whole cell extracts (lanes 8-13) were run on 5% SDS-PAGE and subsequently probed with anti-IRF-3 antibody.

20 (C) 293 cells were co-transfected with myc-tagged CBP expression plasmids (as indicated above the lanes) and IRF-3  $\Delta$ N ( $\Delta$ 9-133) expression plasmid. At 24h after transfection, cells were infected with Sendai virus (lanes 2, 4 and 6) or left uninfected (lanes 1, 3 and 5). Whole cell extracts (300  $\mu$ g) were immunoprecipitated with monoclonal anti-myc-tag antibody 9E10. The immunoprecipitated complexes were run on 5% SDS-PAGE and different forms of IRF-3 in the precipitates were analyzed by immunoblotting with anti-IRF-3 antibody.

(D) Whole cell extracts (30  $\mu$ g) from (C) were also analyzed directly for the expression of myc-tagged CBP proteins by immunoblotting using anti-myc antibody 9E10.

**EXAMPLE****Plasmid constructions and Mutagenesis.**

The IRF-3 expression plasmid was prepared by cloning the *EcoRI*-*XhoI* fragment containing the IRF-3 cDNA from the pSKIRF-3 plasmid downstream of the CMV promoter of CMVBL vector. CMVt-IRF-3 was constructed by cloning of IRF-3 cDNA downstream of the doxycycline-responsive promoter CMVt at the *Bam*HI site of the neo CMVt BL vector (49). cDNAs encoding IRF-3 carboxyl terminal deletion mutations were generated by 28 cycles of PCR amplification with Vent DNA polymerase. DNA oligonucleotide primers were synthesized using an Applied Biosystems DNA/RNA synthesizer. The amino-terminal primer was synthesized with an *EcoRI* restriction enzyme site and the carboxyl-terminal primers were synthesized with *XbaI* restriction enzyme sites at their ends. The PCR products were purified by phenol/chloroform extraction and ethanol precipitation, digested with *EcoRI* and *XbaI*, and inserted into *EcoRI*/*XbaI* sites of CMVBL vector.

The point mutations of IRF-3 were generated by overlap PCR mutagenesis using Vent DNA polymerase. Mutations were confirmed by sequencing. The N-terminal deletion mutations ( $\Delta$ N,  $\Delta$ N2A,  $\Delta$ N3A and  $\Delta$ N5A) of IRF-3 were generated by digestion of the related IRF-3/CMVBL plasmid with *Bam*HI (filled in with Klenow enzyme), partial digestion with *ScaI*, and re-ligation. GFP-IRF-3 expression plasmids were generated by cloning of cDNAs encoding wild type or mutated forms of IRF-3 into the downstream of EGFP in the pEGFP-C1 vector (Clontech). For construction of plasmids encoding

myc-tagged CBP truncated proteins, the cDNAs coding for CBP were generated from the pRC-RSV/mCBP plasmid (provided by Dr. Dimitris Thanos) by PCR amplification. The cDNA fragments were cloned in the downstream of myc-tag in 5' myc-PCDNA3 vector (provided by Dr. Stephane Richard).

#### Generation of IRF-3 cell lines.

Plasmid CMVt-rtTA (49) was introduced into 293 cells by a calcium phosphate-based method. Cells were selected beginning at 48h after transfection for about one week in  $\alpha$ MEM media (GIBCO-BRL) containing 10% heat-inactivated calf serum, glutamine, antibiotics and 2.5 ng/ $\mu$ l puromycin (Sigma). Resistant cells carrying the CMVt-rtTA plasmid (rtTA-293 cells) were then transfected with the CMVt-IRF-3 plasmid. Cells were selected beginning at 48h for a period of approximately 2 weeks in  $\alpha$ MEM containing 10% heat-inactivated calf serum, glutamine, antibiotics, 2.5 ng/ $\mu$ l puromycin and 400  $\mu$ g/ml G418 (Life Technologies, Inc.).

#### Cell culture and transfections.

All transfections for CAT assay were carried out in human embryonic kidney 293 cells or NIH3T3 cells grown in  $\alpha$ MEM (293) or Dulbecco's MEM (NIH3T3) media (GIBCO-BRL) supplemented with 10% calf serum, glutamine and antibiotics. Subconfluent cells were transfected with 5  $\mu$ g of CsCl purified chloramphenicol acetyltransferase (CAT) reporter and expression plasmids by calcium phosphate coprecipitation method (293 cells) or lipofectamine (NIH3T3 cells). The reporter plasmids were the SV $\beta$  CAT and ISG15 CAT reporter

genes (56); also the transfection procedures were previously described (41,56). For individual transfections, 100  $\mu$ g (SV $\beta$  CAT) or 10  $\mu$ g (ISG15 CAT) of total protein extract was assayed for 1-2h at 37°C. The CAT activity was normalized with  $\beta$ -Gal assay. All transfections were performed 3-6 times.

Western blot analysis of IRF-3 modification and degradation.

To characterize the posttranslational regulation of IRF-3 protein, stable or transiently transfected IRF-3 expressing cells were infected with Sendai Virus (80 HAU/ml) or treated with 5 ng/ml TNF- $\alpha$ , either with or without addition of 50  $\mu$ g/ml cycloheximide. In some experiments, cells were treated with either 100  $\mu$ M calpain inhibitor I (ICN), 40  $\mu$ M MG132 proteasome inhibitor, or an equivalent volume of their respective solvent (ethanol) as control. Cells were washed with phosphate-buffered saline and lysed in 10 mM Tris-Cl pH 8.0, 200 mM NaCl, 1 mM EDTA, 1 mM dithiothreitol (DTT), 0.5% Nonidet P-40 (NP-40), 0.5 mM phenylmethanesulfonyl fluoride (PMSF), 5  $\mu$ g/ml leupeptin, 5  $\mu$ g/ml pepstatin, and 5  $\mu$ g/ml aprotinin. Equivalent amounts of whole cell extract (20  $\mu$ g) were subject to SDS-polyacrylamide gel electrophoresis (SDS-PAGE) in a 10% polyacrylamide gel. After electrophoresis, the proteins were transferred to Hybond transfer membrane (Amersham) in a buffer containing 30 mM Tris, 200 mM glycine and 20% methanol for 1h. The membrane was blocked by incubation in phosphate-buffered saline (PBS) containing 5% dried milk for 1h and then probed with IRF-3 antibody in 5% milk/PBS, at a

dilution of 1:3000. These incubations were done at 4°C overnight or at RT for 1-3h. After four 10 minute washes with PBS, membranes were reacted with a peroxidase-conjugated secondary goat anti-rabbit antibody (Amersham) at a dilution of 1:2500. The reaction was then visualized with the enhanced chemiluminescence detection system (ECL) as recommended by the manufacturer (Amersham Corp.).

#### Phosphatase treatment.

Twenty to sixty µg of whole cell extract were treated with 0.3 units of potato acidic phosphatase (Sigma) in a final volume of 30 µl PIPES buffer (10 mM PIPES pH 6.0, 0.5 mM PMSF, 5 µg/ml aprotinin, 1 µg/ml leupeptin, and 1 µg/ml pepstatin) or 5 units of calf intestine alkaline phosphatase (Pharmacia) in 30 µl CIP buffer. The phosphatase inhibitor mix contained 10 mM NaF, 1.5 mM Na<sub>2</sub>MoO<sub>4</sub>, 1 mM β-glycerophosphate, 0.4 mM Na<sub>3</sub>VO<sub>4</sub> and 0.1 µg/ml okadaic acid.

#### Subcellular localization of GFP-IRF-3 proteins.

To analyse the subcellular localization of wild type and mutated forms of IRF-3 proteins in uninfected and virus infected cells, the GFP-IRF-3 expression plasmids (5 µg) were transiently transfected into COS-7 cells by the calcium phosphate coprecipitation method. For virus infection, transfected cells were infected with Sendai virus (80 hemagglutinating units per mL for 2h) at 24h post transfection. GFP fluorescence was analyzed in living cells with a Leica fluorescence microscope using a 40x objective.

#### Electromobility Shift Assay.

Nuclear extracts were prepared from 293 cells at

different times after infection with Sendai virus (80HAU/ml).

In some experiments, extracts were prepared from cells transfected with different IRF-3 expression plasmids, as indicated in individual experiments. Cells were washed in

Buffer A [10 mM HEPES, pH 7.9; 1.5 mM  $MgCl_2$ ; 10 mM KCl; 0.5 mM dithiothreitol (DTT); and 0.5 mM phenylmethylsulfonyl

fluoride (PMSF)] and were resuspended in Buffer A containing 0.1% NP-40. Cells were then chilled on ice for 10 min before centrifugation at 10,000 g. Pellets were then resuspended in

10 Buffer B (20mM HEPES, pH 7.9; 25% glycerol; 0.42 M NaCl; 1.5 mM  $MgCl_2$ ; 0.2 mM EDTA; 0.5 mM DTT; 0.5 mM PMSF; 5  $\mu$ g/ml leupeptin; 5  $\mu$ g/ml pepstatin; 0.5 mM spermidine; 0.15 mM spermine; and 5  $\mu$ g/ml aprotinin). Samples were incubated on ice for 15 min before being centrifuged at 10,000 g. Nuclear extract supernatants were diluted with Buffer C (20 mM HEPES, pH 7.9; 20% glycerol; 0.2 mM EDTA; 50 mM KCl; 0.5 mM DTT; and 0.5 mM PMSF). Nuclear extracts were subjected to EMSA by using a  $^{32}P$ -labelled probe corresponding to the PRDIII region of the IFN- $\beta$  promoter (5'-GGAAACTGAAAGGG-3') or the ISRE

20 region of the ISG-15 promoter

(5'-GATCGGGAAAGGGAAACCGAAACTGAAGCC-3'). The resulting protein-DNA complexes were resolved by 5% polyacrylamide gel and exposed to X-ray film. To demonstrate the specificity of protein-DNA complex formation, 125-fold molar excess of unlabelled oligonucleotide was added to the nuclear extract before adding labelled probe.



Immunoprecipitation and Western analysis of CBP associated proteins.

Whole cell extract (300  $\mu$ g) were prepared from either transfected or untransfected cells and precleared with 5  $\mu$ l of preimmune rabbit serum and 20  $\mu$ l of protein A-Sepharose beads (Pharmacia) for 1 h at 4°C. The extract was incubated with 10  $\mu$ l of anti-CBP antibody A-22 (Santa Cruz) or 2  $\mu$ l anti-myc antibody 9E10 (21) and 30  $\mu$ l of protein A-Sepharose beads for 2-3 h at 4°C. Precipitates were washed 5 times with lysis buffer, eluted by boiling the beads 3 min in 1x SDS sample buffer. Eluted proteins were separated by SDS PAGE, transferred to Hybond transfer membrane. Membranes were incubated with anti-IRF-3 (1:3000) or anti-myc antibody 9E10 (1:1000). Immunocomplexes were detected by using a chemiluminescence-based system.

Virus induced phosphorylation of IRF-3 protein.

IRF-3 is expressed constitutively in various cells and its expression is not enhanced by viral infection or by IFN treatment. To investigate whether the IRF-3 protein is regulated by post-translational modification after virus infection, 293 cells were transiently transfected with an IRF-3 expression plasmid and subsequently infected with Sendai virus 24h later. In cells transfected with CMVBL vector alone, endogenous IRF-3 protein was easily detected using a polyclonal IRF-3 antibody and in cells transfected with the IRF-3 expression plasmid, IRF-3 protein levels were significantly increased (Fig.1, lanes 1 and 3). Interestingly, Sendai virus infection resulted in two

alterations in the expression of IRF-3: 1) an overall decrease in the amount of IRF-3 in transfected and control cells (Fig. 1, lanes 2 and 4) and the generation of a more slowly migrating form of IRF-3 (Fig. 1, compare lanes 1 and 2). In all experiments, the turnover of IRF-3 after virus infection was more pronounced with the endogenous protein than with the transfected proteins (see Fig. 1, as well as others). Because the transfected proteins were driven by the CMV promoter, ongoing synthesis of transfected IRF-3 may partially obscure the turnover of IRF-3.

The kinetics of virus-induced modification of IRF-3 were characterized in a 293 cell line that expressed IRF-3 inducibly under the control of the tetracycline responsive promoter CMVt (25,26). Infection of this cell line (designated rtTA-IRF-3) with Sendai virus resulted in a decrease in the amount of IRF-3 between 12 and 24h after infection (Fig. 2A). Two forms of IRF-3 protein (designated I and II) were detected in uninfected cells (Fig. 2A, lane 1) and following virus infection, a third slowly migrating form of IRF-3 was also detected (Fig. 2A, lanes 4-7). To determine whether the slowest form of IRF-3 was due to virus-induced phosphorylation (P-IRF-3), the different forms of IRF-3 were subjected to treatment *in vitro* with potato acidic phosphatase (PPA) or calf intestine alkaline phosphatase (CIP) and/or phosphatase inhibitors (Fig. 2B). These treatments did not affect the mobilities of forms I and II in uninfected cells (Fig. 2B, lanes 1-3). However, in rtTA-IRF-3 expressing 293 cells infected with Sendai virus

for 12h, an additional slowly migrating, presumably phosphorylated form of IRF-3 was also detected (Fig. 2B, lane 6); this form of IRF-3 completely disappeared following CIP or PPA treatment (Fig. 2B, lanes 6 and 7) but was maintained in the presence of CIP/PPA when phosphatase inhibitors were also added to the reaction (Fig. 2B, lanes 5 and 8).

Mapping the IRF-3 phosphorylation sites.

A series of deletions of IRF-3 were generated to identify the virus-induced phosphorylation site(s) of IRF-3 (Fig. 3A). 293 cells were transiently transfected with IRF-3 deletion mutants and the virus mediated phosphorylation was measured by immunoblotting (Fig. 3B). The results indicated that a virus-induced phosphorylation of IRF-3 occurs at the C-terminal end of IRF-3 since the mutations that contained only the N-terminal part of IRF-3 protein (133, 240, 328, 357 or 394aa) were not phosphorylated (Fig. 3B). Full length and 407aa forms of IRF-3 were phosphorylated as a consequence of virus infection (Fig. 3B, lanes 1-4). C-terminal truncation of IRF-3 to a protein of 394 or 357aa removed the site(s) of inducible phosphorylation (Fig. 3B, lanes 5-8), although the shortened versions of forms I and II were still observed. Also in the IRF-3  $\Delta$ 9-133 mutation ( $\Delta$ N) which had the DNA binding, N-terminal amino acids (aa9 to aa133) removed, both virus induced phosphorylation of IRF-3 and the differential migration of the shortened forms I and II were easily detected (Fig. 3B, lanes 9 and 10). Degradation of the endogenous forms of IRF-3 by virus infection was also detected in this experiment (compare Fig. 3B, lanes 7 and 9

with lanes 8 and 10).

Thus, by deletion analysis, a phosphorylation domain of IRF-3 protein was localized to the region -ISNSHPLSLTSDQ- between amino acids 395 and 407. Point

mutations in the several putative Ser and Thr phosphorylation residues within this region were generated in the full length protein and the  $\Delta 9-133$  ( $\Delta N$ ) protein (Fig. 4A). In the IRF-3 cDNA encoding these proteins, the

Ser-396/Ser398/Ser-402/Thr-404/Ser-405 residues were replaced by alanine (5A), as were the three residues

Ser-402/Thr-404/Ser-405 (3A) and the two residues

Ser-396/Ser-398 (2A). Transfection of these plasmids into 293 cells and subsequent virus infection revealed that full length wild type IRF-3 was phosphorylated (Fig. 4B, lanes 4 and 8), whereas the IRF-3 proteins containing 2A and 5A

mutations were no longer phosphorylated in virus infected cells (Fig. 4B, lanes 6 and 10). Interestingly, IRF-3-3A was also very weakly phosphorylated as a consequence of virus infection, thus implicating Ser-402/Thr-404/Ser-405 as

potential secondary sites of phosphorylation. Using the  $\Delta N$  IRF-3 protein and the relevant point mutations,

phosphorylation was detected with  $\Delta N$  (Fig. 4B, lane 12) but not with  $\Delta N$ -2A and  $\Delta N$ -5A (Fig. 4B, lanes 14 and 18);

likewise,  $\Delta N$ -3A displayed very weak phosphorylation (Fig. 4B, lane 16).

These experiments thus implicate Ser-396 and Ser-398 as critical sites of virus-induced phosphorylation of IRF-3; however, Ser-402/Thr-404/Ser-405 residues also

---contribute to the observed phosphorylation, since the migration of phosphorylated  $\Delta N$ -3A is significantly faster than  $\Delta N$  and the phosphorylation level is decreased (Fig. 4B, lanes 12 and 16). Another study suggested the involvement of the Ser residues at aa385 and 386 as potential phosphoacceptor sites (67); in studies with the S385A/S386A mutation, applicant found no evidence for inducible phosphorylation at these sites. However, since these sites represent consensus sites for CKI and CKII, constitutive phosphorylation is a possibility.

IRF-3 phosphorylation induces cytoplasmic to nuclear translocation of IRF-3.

Initial studies indicated that IRF-3 was localized in the cytoplasm of uninfected cells (67); to investigate the role of phosphorylation on IRF-3 localization, wild type and point mutated forms of IRF-3 were linked to green fluorescent protein (GFP), transfected into COS-7 cells and examined for Sendai virus induced changes in subcellular localization (Fig. 5). In uninfected cells, GFP-IRF-3 localized exclusively to the cytoplasm; Sendai virus infection resulted in translocation of IRF-3 to the nucleus within 8h in 90-95% of the cells (Fig. 5A and B). Mutation of the Ser/Thr cluster in GFP-IRF-3(5A) completely abrogated virus-induced cytoplasmic to nuclear translocation (Fig. 5, C and D). Interestingly, the substitution of the Ser/Thr cluster with the phosphomimetic Asp in GFP-IRF-3(5D) likewise altered subcellular localization. IRF-3(5D) localized both to the nucleus and cytoplasm in uninfected cells (Fig. 5E), while

~~virus infection resulted in an intense nuclear pattern of~~  
 IRF-3(5D) fluorescence (Fig. 5F). Point mutation of a  
 putative nuclear export signal in IRF-3, the L145A/L146A  
 modification - termed IRF-3(NES) - also changed subcellular  
 localization of IRF-3. In uninfected cells, GFP-IRF-3(NES)  
 was localized to the nucleus and cytoplasm, with a  
 homogeneous, extra-nucleolar pattern of nuclear staining.  
 After virus infection, GFP-IRF-3(NES) localized to the  
 nucleus with an intense speckled pattern of nuclear  
 10 fluorescence in greater than 95% of the cells, suggesting  
 that IRF-3(NES) may be trapped in the nucleus associated with  
 the nuclear pore complex.

Transactivation of PRDI/PRDIII and ISRE promoters by IRF-3.

Next, the capacity of IRF-3 to regulate gene  
 expression was analysed by transient transfection in human  
 293 and murine NIH3T3 cells using the IFN $\beta$  and ISG-15  
 promoters in reporter gene assays. Expression of NF- $\kappa$ B  
 RelA(p65), IRF-1 and IRF-3 alone minimally induced IFN $\beta$   
 promoter activity between 3 to 4 fold (Fig. 6A and B), as  
 20 shown previously (24,56,61). Introduction of the C-terminal  
 point mutants - IRF-3(2A), IRF-3(3A) IRF-3(5A) - reduced the  
 low transactivation capacity of IRF-3 to control levels (Fig.  
 6A). Interestingly, deletion of the C-terminal 20aa of IRF-3  
 to IRF-3(407) stimulated IFN $\beta$  activity about 6 fold,  
 indicative of the removal of an inhibitory domain in IRF-3.  
 However, further deletion to 394, 357 or 240 abrogated  
 transactivation potential (Fig. 6A). Mutation of the NES  
 element was not sufficient to stimulate IFN $\beta$  activity.

~~Strikingly, the substitution of the Ser/Thr cluster at~~  
aa397-405 in IRF-3 with the phosphomimetic Asp generated a  
very strong, constitutive transactivator protein that alone  
stimulated the IFN $\beta$  promoter 90 fold.

As shown previously, high level induction of the  
IFN $\beta$  promoter requires synergistic activation by NF- $\kappa$ B and  
IRF proteins (24,61). To analyse the properties of IRF-3 in  
synergistic activation of the IFN $\beta$  promoter, co-expression  
studies were performed using RelA(p65) expression plasmid and  
different wild type and mutant forms of IRF-3 (Fig. 6B).

Co-expression of RelA and IRF-1 or RelA and IRF-3 stimulated  
IFN $\beta$ -CAT activity by 20-25 fold. IRF-3(407) and RelA(p65)  
stimulated IFN $\beta$  activity about 40 fold, supporting the idea  
of the removal of an inhibitory domain in IRF-3, whereas both  
the IRF-3(394) and the IRF-3(NES) failed to synergise with  
RelA in the activation of the IFN $\beta$  promoter. RelA and  
IRF-3(NES) produced a relatively weak 8 fold induction of  
IFN $\beta$  expression, indicating that nuclear localization is not  
sufficient for IRF-3 activation. The combination of RelA and  
IRF-3(5D) produced an 80 fold stimulation of IFN $\beta$  promoter  
activity (Fig. 6B); together with the above data, IRF-3(5D)  
alone appears to be capable of full stimulation of the IFN $\beta$   
promoter and further synergy with RelA is not observed  
(compare Fig. 6A and B). Surprisingly, IRF-3(5A) and RelA  
produced a 30 fold stimulation, suggesting that 5A can still  
synergise with RelA, despite mutation of the Ser/Thr cluster.

The transactivation potential of IRF-3 was also  
analysed using the ISG-15 promoter, an ISRE containing

regulatory element (Fig. 6C). As shown previously (2), and above for the IFN $\beta$  promoter, IRF-3 alone weakly activated the ISG-15 promoter; in the context of this regulatory element, IRF-3 was weaker than IRF-1, which produced a 9 fold

stimulation. Again deletion of the C-terminal 20aa of IRF-3 generated a protein that stimulated gene expression; with the ISG-15 promoter, a 12 fold induction was observed; IRF-3(394) and IRF-3(357) did not stimulate gene expression but rather slightly repressed ISG-15. Again remarkably, IRF-3(5D)

10 produced a 50 fold induction of the ISG-15 promoter (Fig. 6C), thus demonstrating that substitution of the Ser/Thr sites with the phosphomimetic Asp generated a constitutively active form of IRF-3 that functioned as a very strong activator of promoters containing PRDI/PRDIII or ISRE regulatory elements.

#### Inhibition of IRF-3 degradation.

Another consequence of virus infection is the degradation of the IRF-3. Since phosphorylation of proteins is functionally associated with the process of protein  
20 degradation via the ubiquitin-dependent proteasome pathway (53,57,60), the effect of proteasome inhibitors on virus-induced turnover of IRF-3 was examined. In cells transfected with the  $\Delta$ N and  $\Delta$ N5A forms of IRF-3, virus induced degradation of full length (endogenous) forms of IRF-3 (Fig. 7A, lanes 1 and 4) and the truncated  $\Delta$ N (Fig. 7B, lanes 1 and 4) was detected. Addition of the protease inhibitor calpain inhibitor I or the proteasome inhibitor MG132 blocked virus-induced IRF-3 degradation (Fig. 7A and



7B, lanes 4-6). Particularly with the  $\Delta N$  protein, the accumulation of the phosphorylated form of  $\Delta N$  was also detected in virus infected cells (Fig. 7B, lanes 5 and 6), suggesting that phosphorylation of IRF-3 may represent a signal for subsequent degradation by the proteasome pathway. To confirm this idea, the 5A point mutated form of IRF-3 was analysed; the IRF-3- $\Delta N5A$  protein was resistant to virus induced degradation (Fig. 7C, lanes 1 and 4); no further stabilization of IRF-3- $\Delta N5A$  occurred with calpain inhibitor I or MG132 addition and no phosphorylated IRF-3 was detected (Fig. 7C, lanes 4-6). These experiments demonstrate that virus dependent phosphorylation of the C-terminal of IRF-3 represents a signal for subsequent proteasome mediated degradation.

#### Interaction between IRF-3 and CBP in virus infected cells.

To examine the possibility that IRF-3 associated with the co-activator CBP/p300 (Fig. 8A) following Sendai virus infection, CBP was immunoprecipitated from virus-infected cells with anti-CBP antibody; an immunoblot for IRF-3 revealed that IRF-3 was co-precipitated from virus-infected cells but not from uninfected cells (Fig. 8B, lanes 2 and 3). This interaction was observed clearly in cells co-transfected with the IRF-3 expression plasmid (Fig. 8B, lane 3 ) but was not seen when the immunoprecipitation was performed with pre-immune serum (Fig. 8B, lane 7). The endogenous IRF-3 also co-precipitated from virus-infected cells (Fig. 8B, lane 1). However, mutation of the Ser/Thr residues identified as the virus inducible phosphorylation

sites abrogated the association of IRF-3 with CBP. In particular, IRF-3(2A) and IRF-3(5A) were detected in whole cell extract immunoblot but not in the CBP immunoprecipitate (Fig. 8B, compare lanes 4 and 6 with lanes 11 and 13). With the IRF-3(3A) mutant, interaction with CBP was still observed (Fig. 8B, lane 5). The high background in all lanes represents secondary antibody reactivity with rabbit IgG from the immunoprecipitation. Immunoblot analysis of the whole cell extracts revealed that phosphorylated IRF-3, as well as forms I and II were present in virus infected cells (Fig. 8B, lane 10) and in cells transfected with 2A, 3A and 5A the forms I and II were observed but not the phosphorylated form of IRF-3 (Fig. 8B, lanes 11-13).

CBP has several domains that bind transcription factors, designated CBP1, CBP2, and CBP3 respectively (Fig. 8A, reviewed in (28)). To determine which domain of CBP interacts with IRF-3, the three specific subdomains were myc-tagged at the 5' end by subcloning into the pCDNA3 vector (Fig. 8A). 293 cells were co-transfected with these myc-tagged CBP expression plasmids together with the IRF-3  $\Delta$ N ( $\Delta$ 9-133) expression plasmid. At 24h after transfection, cells were infected with Sendai virus, co-immunoprecipitated with anti-myc antibody 16h later (21) and then immunoblotted for IRF-3. Endogenous IRF-3 and transfected IRF-3  $\Delta$ N proteins co-precipitated with CBP-3 from virus-infected cells but not from uninfected cells (Fig. 8C, lane 6). In cells co-transfected with CBP-1 and CBP-2, no endogenous or transfected  $\Delta$ N IRF-3 was detected (Fig. 8C, lanes 1-4).

- Immunoblot analysis of the whole-cell extracts revealed that all three myc-tagged CBP proteins were efficiently expressed in uninfected and virus infected cells (Fig. 8D). These results demonstrate that IRF-3 binds to the C-terminal domain of CBP in virus infected cells and interaction with CBP requires Ser-396/Ser-398 phosphorylation of IRF-3 but not at Ser-402/Thr-404/Ser-405.

#### REFERENCES

1. Arany, Z., Sellers, W.R., Livingston, D.M. and Eckner, R.  
10 1994. E1A-associated p300 and CREB-associated CBP belong to a conserved family of coactivators. Cell 77:799-800.
2. Au, W.-C., Moore, P.A., Lowther, W., Juang, Y.-T. and Pitha, P.M. 1995. Identification of a member of the interferon regulatory factor family that binds to the interferon-stimulated response element and activates expression of interferon-induced genes. Proc.Natl.Acad.Sci.USA 92:11657-11661.
3. Avantaggiati, M.L., Ogryzko, V., Gardner, K., Giordano, A., Levine, A.S. and Kelly, K. 1997. Recruitment of  
20 p300/CBP in p53-dependent signal pathways. Cell 89:1175-1184.
4. Baldwin, A.S.Jr. 1996. The NF- $\kappa$ B and I $\kappa$ B proteins: new discoveries and insights. Annu.Rev.Immunol. 14:649-681.
5. Bannister, A.J. and Kouzarides, T. 1996. The CBP coactivator is a histone acetyltransferase. Nature 384:641-643.
6. Beauparlant, P. and Hiscott, J. 1996. Biological and biochemical inhibitors of the NF- $\kappa$ B/Rel proteins and cytokine

synthesis. CytGrowthFactRev 7:175-190.

7. Bhattacharya, S., Eckner, R., Grossman, S., Oldread, E., Arany, Z., D'Andrea, A. and Livingston, D.M. 1996.

Cooperation of Stat2 and p300/CBP by interferon- $\alpha$ . Nature 383:344-347.

8. Bluysen, H.A.R., Durbin, J.E. and Levy, D.E. 1996. ISGF3 $\gamma$  p48, a specificity switch for interferon activated transcription factors. CytGrowthFactRev 7:11-17.

9. Bovolenta, C., Driggers, P.H., Marks, M.S., Medin, J.A., Politis, A.D., Vogel, S.N., Levy, D.E., Sakaguchi, K., Appella, E., Coligan, J.E. and Ozato, K. 1994. Molecular interactions between interferon consensus sequence binding protein and members of the interferon regulatory factor family. Proc.Natl.Acad.Sci.USA 91:5046-5050.

10. Bragança, J., Génin, P., Bandu, M.-T., Darracq, N., Vignal, M., Cassé, C., Doly, J. and Civas, A. 1997. Synergism between multiple virus-induced-factor-binding elements involved in the differential expression of IFN-A genes. J.Biol.Chem. 272: 22154-22162.

20 11. Brass, A.L., Kehrl, E., Eisenbeis, C.F., Storb, U. and Singh, H. 1996. Pip, a lymphoid-restricted IRF, contains a regulatory domain that is important for autoinhibition and ternary complex formation with the Ets factor PU.1. Genes Dev. 10:2335-2347.

12. Chen, H., Lin, R.J., Schiltz, R.L., Chakravarti, D., Nash, A., Nagy, L., Privalsky, M.L., Nakatani, Y. and Evans, R.M. 1997. Nuclear receptor coactivator ACTR is a novel histone acetyltransferase and forms a multimeric activation

~~complex with P/CAF and CBP/p300. Cell 90:569-580.~~

13. Chrivia, J.C., Kwok, R.P.S., Lamb, N., Hagiwara, M., Montminy, M.R. and Goodman, R.H. 1993. Phosphorylated CREB binds specifically to the nuclear protein CBP. Nature

365:855-859.

14. Cohen, L. and Hiscott, J. 1992. Characterization of TH3, an induction specific protein interacting with the interferon- $\beta$  promoter. Virol. 191:589-599.

15. Crepieux, P., Coll, J. and Stehelin, D. 1994. The Ets family of proteins: weak modulators of gene expression in quest for transcriptional partners. CritRevOncogen 5:615-638.

16. Daly, C. and Reich, N.C. 1993. Double-stranded RNA activates novel factors that bind to the interferon stimulated response element. Mol.Cell.Biol. 13:3756-3764.

17. Darnell Jr., J.E., Kerr, I.M. and Stark, G.R. 1994. Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. Science 264:1415-1421.

18. Driggers, P.H., Ennist, D.L., Gleason, S.L., Mak, W.-H., Marks, M.S., Levi, B.-Z., Flanagan, J.R., Appella, E. and Ozato, K. 1990. An interferon  $\gamma$ -regulated protein that binds the interferon-inducible enhancer element of major histocompatibility complex class I genes.

Proc.Natl.Acad.Sci.USA 87:3743-3747.

19. Eisenbeis, C.F., Singh, H. and Storb, U. 1995. Pip, a novel IRF family member, is a lymphoid-specific, PU.1-dependent transcriptional activator. Genes Dev.

9:1377-1387.

20. Escalante, C.R., Yie, J., Thanos, D. and Aggarwal, A.K. 1998. Structure of IRF-1 with bound DNA reveals determinants of interferon regulation. *Nature* 391:103-106.

21. Evan, G.I. and Bishop, J.M. 1985. Isolation of monoclonal antibodies specific for the human c-myc proto-oncogene product. *Mol.Cell.Biol.* 4:2843-2850.

22. Fujita, T., Kimura, Y., Miyamoto, M., Barsoumian, E.L. and Taniguchi, T. 1989. Induction of endogenous IFN- $\alpha$  and IFN- $\beta$  genes by a regulatory transcription factor IRF-1. *Nature* 337:270-272.

23. Fujita, T., Sakakibara, J., Sudo, Y., Miyamoto, M., Kimura, Y. and Taniguchi, T. 1988. Evidence for a nuclear factor(s), IRF-1, mediating induction and silencing properties to human IFN- $\beta$  gene regulatory elements. *EMBO J.* 7:3397-3405.

24. Garoufalidis, E., Kwan, I., Lin, R., Mustafa, A., Pepin, N., Roulston, A., Lacoste, J. and Hiscott, J. 1994. Viral induction of the human interferon  $\beta$  promoter: modulation of transcription by NF- $\kappa$ B/rel proteins and interferon regulatory factors. *J.Virol.* 68:4707-4715.

25. Gossen, M. and Bujard, H. 1992. Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proc.Natl.Acad.Sci.USA* 89:5547-5551.

26. Gossen, M., Freundlieb, S., Bender, G., Müller, G., Hillen, W. and Bujard, H. 1995. Transcriptional activation by tetracyclines in mammalian cells. *Science* 268:1766-1769.

27. Gu, W. and Roeder, R.G. 1997. Activation of p53 sequence-specific DNA binding by acetylation of p53 C-terminal domain. *Cell* 90:595-606.
28. Gu, W., Shi, X.L. and Roeder, R.G. 1997. Synergistic activation of transcription by CBP and p53. *Nature* 387:819-823.
29. Génin, P., Bragança, J., Darracq, N., Doly, J. and Civas, A. 1995. A novel PRDI and TG binding activity involved in virus-induced transcription of IFN-A genes. *NuclAcidRes* 23:5055-5063.
30. Harada, H., Fujita, T., Miyamoto, M., Kimura, Y., Maruyama, M., Furia, A., Miyata, T. and Taniguchi, T. 1989. Structurally similar but functionally distinct factors, IRF-1 and IRF-2, bind to the same regulatory elements of IFN and IFN-inducible genes. *Cell* 58:729-739.
31. Harada, H., Matsumoto, M., Sato, M., Kashiwazaki, Y., Kimura, T., Kitagawa, M., Yokochi, T., Tan, R.S.-P., Takasugi, T., Kadokawa, Y., Schindler, C., Schreiber, R.D., Noguchi, S. and Taniguchi, T. 1996. Regulation of IFN- $\alpha/\beta$  genes: evidence for a dual function of the transcription factor complex ISGF3 in the production and action of IFN- $\alpha/\beta$ . *GenestoCells* 1:995-1005.
32. Harada, H., Willison, K., Sakakibara, J., Miyamoto, M., Fujita, T. and Taniguchi, T. 1990. Absence of type I IFN system in EC cells: transcriptional activator (IRF-1) and repressor (IRF-2) genes are developmentally regulated. *Cell* 63:903-913.
33. Hiscott, J., Nguyen, H. and Lin, R. 1995. Molecular

mechanisms of interferon  $\beta$  gene induction. *SeminVirol*  
6:161-173.

34. Holtschke, T., Löhler, J., Kanno, Y., Fehr, T., Giese, N., Rosenbauer, F., Lou, J., Knobloch, K.-P., Gabriele, L.,

Waring, J.F., Bachmann, M.F., Zingernagel, R.M., Morse III, H.C., Ozato, K. and Horak, I. 1996. Immunodeficiency and chronic myelogenous leukemia-like syndrome in mice with a targeted mutation of the ICSBP gene. *Cell* 87:307-317.

35. Ihle, J.N. 1996. STATs: signal transducers and  
10 activators of transcription. *Cell* 84:331-334.

36. Kawakami, T., Matsumoto, M., Sato, M., Harada, H., Taniguchi, T. and Kitigawa, M. 1995. Possible involvement of the transcription factor ISGF3 $\gamma$  in virus-induced expression of the IFN- $\beta$  gene. *FEBS Lett.* 358:225-229.

37. Kim, T.K. and Maniatis, T. 1998. The mechanism of transcriptional synergy of an in vitro assembled interferon  $\beta$  enhanceosome. *Mol.Cell* 1:119-129.

38. Kimura, T., Kadokawa, Y., Harada, H., Matsumoto, M., Sato, M., Kashiwazaki, Y., Tarutani, M., Tan, R.S-P.,

20 Takasugi, T., Matsuyama, T., Mak, T.M., Noguchi, S. and Taniguchi, T. 1996. Essential and non-redundant roles of p48 (ISGF3 $\gamma$ ) and IRF-1 in both type I and type II interferon responses, as revealed by gene targeting studies. *GenestoCells* 1:115-124.

39. Levy, D.E. 1995. Interferon induction of gene expression through the Jak-Stat pathway. *SeminVirol* 6:181-190.

40. Lill, N.L., Grossman, S.R., Ginsberg, D., DeCaprio, J.



and Livingston, D.M. 1997. Binding and modulation of p53 by p300/CBP coactivators. *Nature* 387:823-827.

41. Lin, R., Beauparlant, P., Makris, C., Meloche, S. and Hiscott, J. 1996. Phosphorylation of I $\kappa$ B $\alpha$  in the C-terminal

PEST domain by casein kinase II affects intrinsic protein stability. *Mol.Cell.Biol.* 16:1401-1409.

42. Lin, R., Mustafa, A., Nguyen, H. and Hiscott, J. 1994. Mutational analysis of interferon (IFN) regulatory factors 1 and 2: Effects on the induction of IFN- $\beta$  gene expression.

10 *J.Biol.Chem.* 269:17542-17549.

43. Matsuyama, T., Grossman, A., Mittrfcker, H.-W., Siderovski, D.P., Kiefer, F., Kawakami, T., Richardson, C.D., Taniguchi, T., Yoshinaga, S.K. and Mak, T.W. 1995.

Molecular cloning of LSIRF, a lymphoid-specific member of the interferon regulatory factor family that binds the interferon-stimulated response element (ISRE). *NuclAcidRes* 23:2127-2136.

44. Matsuyama, T., Kimura, T., Kitagawa, M., Watanabe, N., Kundig, T., Amakawa, R., Kishihara, K., Wakeham, A., Potter,

20 J., Furlonger, C., Narendran, A., Suzuki, H., Ohashi, P., Paige, C., Taniguchi, T. and Mak, T. 1993. Targeted disruption of IRF-1 or IRF-2 results in abnormal type I IFN induction and aberrant lymphocyte development. *Cell* 75:83-97.

45. Merika, M., Williams, A., Chen, G., Collins, T. and Thanos, D. 1998. Recruitment of CBP/p300 by the IFN $\beta$  enhanceosome is required for synergistic activation of transcription. *Mol.Cell* 1:277-287.

46. Mittrücker, H.-W., Matsuyama, T., Grossman, A., Kündig, T.M., Potter, J., Shahinian, A., Wakeham, A., Patterson, B., Ohashi, P.S. and Mak, T.W. 1997. Requirement for the transcription factor LSIRF/IRF4 for mature B and T lymphocyte function. *Science* 275:540-543.

47. Miyamoto, M., Fujita, T., Kimura, Y., Maruyama, M., Harada, H., Sudo, Y., Miyata, T. and Taniguchi, T. 1988. Regulated expression of a gene encoding a nuclear factor, IRF-1, that specifically binds to the IFN- $\beta$  gene regulatory elements. *Cell* 54:903-913.

48. Nguyen, H., Hiscott, J. and Pitha, P.M. 1997. The growing family of IRF transcription factors. *CytGrowthFactRev* 8:in press.

49. Nguyen, H., Lin, R. and Hiscott, J. 1997. Activation of multiple growth regulatory genes following inducible expression of IRF-1 or IRF/RelA fusion proteins. *Oncogene* 15:1425-1435.

50. Ogryzko, V.V., Schiltz, R.L., Russanova, V., Howard, B.H. and Nakatani, Y. 1996. The transcriptional coactivators p300 and CBP are histone acetyltransferases. *Cell* 87:953-959.

51. Palombella, V. and Maniatis, T. 1992. Inducible processing of interferon regulatory factor-2. *Mol.Cell.Biol.* 12:3325-3336.

52. Pitha, P.M. and Au, W.-C. 1995. Induction of interferon  $\alpha$  gene expression. *SeminVirol* 6:151-159.

53. Read, M.A., Neish, A.S., Luscinskas, F.W., Palombella, V.J., Maniatis, T. and Collins, T. 1995. The proteasome

~~pathway is required for cytokine-induced~~

endothelial-leukocyte adhesion molecule expression. *Immunity* 2:493-506.

54. Reis, L.F.L., Harada, H., Wolchok, J.D., Taniguchi, T.

and Vilcek, J. 1992. Critical role of a common transcription factor, IRF-1, in the regulation of IFN- $\beta$  and IFN-inducible genes. *EMBO J.* 11:185-193.

55. Russo, J.J., Bohenzky, R.A., Chien, M.-C., Chen, J., Yan, M., Maddalena, D., Parry, J.P., Peruzzi, D., Edelman, I.S.,

10 Chang, Y. and Moore, P. 1996. Nucleotide sequence of the kaposi sarcoma-associated herpesvirus (HHV8).

*Proc.Natl.Acad.Sci.USA* 93:14862-14867.

56. Schafer, S., Lin, R., Moore, P., Hiscott, J. and Pitha, P.M. 1998. Regulation of type 1 interferon gene expression by interferon regulatory factor 3. *J.Biol.Chem.*

273:2714-2720.

57. Scherer, D.C., Brockman, J.A., Chen, Z., Maniatis, T. and Ballard, D.W. 1995. Signal-induced degradation of I $\kappa$ B $\alpha$  requires site-specific ubiquitination.

20 *Proc.Natl.Acad.Sci.USA* 92:11259-11263.

58. Schindler, C. and Darnell Jr., J.E. 1995.

Transcriptional responses to polypeptide ligands: the JAK-STAT pathway. *Ann.Rev.Biochem.* 64:621-651.

59. Sharf, R., Meraro, D., Azriel, A., Thornton, A.M., Ozato, K., Petricoin, E.F., Larner, A.C., Schaper, F., Hauser, H.

and Levi, B.-Z. 1997. Phosphorylation events modulate the ability of interferon consensus sequence binding protein to interact with interferon regulatory factors and to bind DNA.

J.Biol.Chem. 272:9785-9792.

60. Thanos, D. and Maniatis, T. 1995. NF- $\kappa$ B: a lesson in family values. Cell 80:529-532.

61. Thanos, D. and Maniatis, T. 1995. Identification of the rel family members required for virus induction of the human  $\beta$  interferon gene. MolCellBiol 15:152-164.

62. Veals, S.A., Schindler, C., Leonard, D., Fu, X.-Y., Aebersold, R., Darnell Jr., J.E. and Levy, D.E. 1992.

10 Subunit of an  $\alpha$ -interferon-responsive transcription factor is related to interferon regulatory factor and myb families of DNA-binding proteins. MolCellBiol 12:3315-3324.

63. Vilcek, J. and Sen, G. Interferons and other cytokines. In: Virology, edited by Fields, B., Knipe, D.M. and Howley, P.M. Philadelphia: Lippincott-Raven, 1996, p. 375-399.

64. Weisz, A., Marx, P., Sharf, R., Appella, E., Driggers, P.H., Ozato, K. and Levi, B.-Z. 1992. Human interferon consensus sequence binding protein is a negative regulator of enhancer elements common to interferon-inducible genes.

J.Biol.Chem. 267:25589-25596.

20 65. Whiteside, S.T., King, P. and Goodbourn, S. 1994. A truncated form of the IRF-2 transcription factor has the properties of a postinduction repressor of interferon- $\beta$  gene expression. J.Biol.Chem. 269:27059-27065.

66. Yamagata, T., Nishida, J., Tanaka, T., Sakai, R., Mitani, K., Yoshida, M., Taniguchi, T., Yazaki, Y. and Hirai, H. 1996. A novel interferon regulatory factor family transcription factor, ICSAT/Pip/LSIRF, that negatively regulates the activity of interferon-regulated genes.

MolCellBiol 16:1283-1294.

67. Yoneyama, M., Suhara, W., Fukuhara, Y. and Fujita, T.  
1997. Direct activation of a factor complex composed of  
IRF-3 and CBP/p300 by virus infection. J.Interferon Cytokine

---

Res. 17:S53.

68. Zhang, J.J., Vinkemeier, U., Gu, W., Chakravarti, D.,  
Horvath, C.M. and Darnell, J.E. 1996. Two contact regions  
between STAT1 and CBP/p300 in interferon  $\gamma$  signalling.  
Proc.Natl.Acad.Sci.USA 93:15092-15096.

10 69. Zhang, L. and Pagano, J.S. 1997. IRF-7, a new  
interferon regulatory factor associated with Epstein Barr  
Virus latency. Mol.Cell.Biol. 17:5748-5757.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE  
PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A phosphorylated form of IRF-3 wherein Ser-396 and Ser-398 residues are phosphorylated.
2. IRF-3 according to claim 1 wherein one or more of Ser-402, Thr-404 and Ser-405 are phosphorylated.
3. A mutant form of IRF-3 wherein one or more of Ser-396, Ser-398, Ser-402, Thr-404 and Ser-405 are substituted by phosphomimetic Asp or phosphomimetic Glu.
4. Use of the IRF-3 according to any one of claims 1 through 3 to activate one or more genes selected from interferon genes and cytokine genes.
5. Use of the IRF-3 according to any one of claims 1 through 3 to treat virus infections.

Smart & Eickar  
Ottawa, Canada  
Patent Agents

Figure 1






Figure 2

**A****B**



A




Figure 3B

B




Figure 4A

A






Figure 4B


B



WIRF-3



B WIRF-3




C IRF-3(5A)




D IRF-3(5A)




E IRF-3(5E)



F IRF-3(5E)



G IRF-3(NES)



H IRF-3(NES)




Figure 2A




Figure 6B


**B. IFN $\beta$ -CAT**

Figure 6C






Figure 7




Figure 8A




Figure 8B

B




Figure 8C

